

[First Hit](#) [Previous Doc](#) [Next Doc](#) [Go to Doc#](#)

☐ [Generate Collection](#) [Print](#)

L8: Entry 3 of 24

File: PGPB

Jan 29, 2004

PGPUB-DOCUMENT-NUMBER: 20040018235
PGPUB-FILING-TYPE: new
DOCUMENT-IDENTIFIER: US 20040018235 A1

TITLE: Controlled release pharmaceutical composition

PUBLICATION-DATE: January 29, 2004

INVENTOR-INFORMATION:

NAME	CITY	STATE	COUNTRY
Tanizawa, Yoshio	Fuji-shi		JP
Shimokawa, Tatsuharu	Fuji-shi		JP
Ogawa, Hirotada	Fuji-shi		JP
Watanabe, Mayumi	Fuji-shi		JP
Ohashi, Chihiro	Fuji-shi		JP
Kawashima, Hiroyuki	Fuji-shi		JP
Shinoda, Yasuo	Shizuoka-shi		JP
Inagi, Toshio	Mishima-shi		JP

ASSIGNEE-INFORMATION:

NAME	CITY	STATE	COUNTRY	TYPE	CODE
KOWA CO., LTD	Nagoya-shi		JP	03	
Nissan Chemical Industries Limited	Tokyo		JP	03	

APPL-NO: 10/461432 [PALM]

DATE FILED: June 16, 2003

RELATED-US-APPL-DATA:

Application is a non-provisional-of-provisional application 60/388740, filed June 17, 2002,

INT-CL-PUBLISHED: [07] A61K 31/366, A61K 9/22, A61K 31/225

INT-CL-CURRENT:

TYPE	IPC	DATE
CIPS	<u>A61 K 9/20</u>	20060101
CIPS	<u>A61 K 9/50</u>	20060101
CIPS	<u>A61 K 31/47</u>	20060101
CIPS	<u>A61 K 9/24</u>	20060101

US-CL-PUBLISHED: 424/468; 514/460, 514/548

US-CL-CURRENT: 424/468; 514/460, 514/548

REPRESENTATIVE-FIGURES: NONE

ABSTRACT:

The present invention is drawn to a pharmaceutical composition characterized by containing a composition (A) which contains pitavastatin, a salt thereof, or an ester thereof and which initiates release thereof at least in the stomach, and an enteric composition (B) which contains pitavastatin, a salt thereof, or an ester thereof. By use of the controlled release pharmaceutical composition of the present invention, the blood level of pitavastatin can be maintained at an appropriate level immediately after administration over a long period of time. Thus, highly safety and effective treatment of hypercholesterolemia can be performed.

[Previous Doc](#)[Next Doc](#)[Go to Doc#](#)

[First Hit](#)[Previous Doc](#)[Next Doc](#)[Go to Doc#](#)[Generate Collection](#)[Print](#)

L8: Entry 3 of 24

File: PGPB

Jan 29, 2004

DOCUMENT-IDENTIFIER: US 20040018235 A1

TITLE: Controlled release pharmaceutical composition

Abstract Paragraph:

The present invention is drawn to a pharmaceutical composition characterized by containing a composition (A) which contains pitavastatin, a salt thereof, or an ester thereof and which initiates release thereof at least in the stomach, and an enteric composition (B) which contains pitavastatin, a salt thereof, or an ester thereof. By use of the controlled release pharmaceutical composition of the present invention, the blood level of pitavastatin can be maintained at an appropriate level immediately after administration over a long period of time. Thus, highly safety and effective treatment of hypercholesterolemia can be performed.

Application Filing Date:

20030616

Summary of Invention Paragraph:

[0001] The present invention relates to a controlled release pharmaceutical composition containing pitavastatin, which is an HMG-CoA reductase inhibitor, a salt thereof, or an ester thereof.

Summary of Invention Paragraph:

[0002] Pitavastatin and salts and esters thereof are known to be useful as drugs for treating hypercholesterolemia, by virtue of their excellent HMG-CoA reductase inhibitory activity (U.S. Pat. No. 5,856,336 and EP 0,304,063). Hypercholesterolemia treatment drugs such as the pitavastatin-containing drugs are used in the form of peroral preparations such as tablets, granules and capsules (U.S. Pat. No. 6,465,477 and WO 97/23200). In general, peroral drug preparations are designed such that the blood level of the active ingredient reaches its peak 0.5 to 3 hours after ingestion, and the level rapidly decreases afterward. However, since cholesterol is synthesized in the body during the period from midnight to morning, it is highly likely that blood level of the active ingredient does not match the time at which biosynthesis of cholesterol occurs.

Summary of Invention Paragraph:

[0003] Although pitavastatin and salts and esters thereof are also known to exert high activity and have high safety, an excessively high blood level thereof is not preferred, from the viewpoint of prevention against the side effects.

Summary of Invention Paragraph:

[0004] In addition, it is also desired for pitavastatin and salts and esters thereof to maintain an excellent hypercholesterol-reducing action for a long period of time.

Summary of Invention Paragraph:

[0005] In view of the foregoing, an object of the present invention is to provide a controlled release pharmaceutical composition which can reliably maintain an appropriate blood level of pitavastatin, a salt thereof, or an ester thereof over a long period of time.

Summary of Invention Paragraph:

[0006] The present inventors have carried out extensive studies in order to develop a controlled release pharmaceutical composition containing pitavastatin, a salt thereof, or an ester thereof. A first approach to resolution of the aforementioned problems is a sustained release drug preparation exhibiting zero-order drug release. However, when such a sustained release drug preparation exhibiting zero-order drug release is administered singly, increasing the blood level thereof immediately after administration is difficult, resulting in a considerably low level of the maximum blood concentration (C.sub.max) as compared with ordinary drug preparations of the same dose. In addition, the area under the blood concentration curve (AUC)--an index of hypercholesterol-reducing action--decreases considerably. Thus, the first approach fails to resolve the aforementioned problems.

Summary of Invention Paragraph:

[0007] Consequently, the present inventors have studied the absorption kinetics of pitavastatin and salts and esters thereof, and have found that pitavastatin and salts and esters thereof are absorbed most effectively in the duodenum and also effectively in the large intestine as well as the small intestine, which contrasts greatly with the general tendency for ordinary drugs to be absorbed in the small intestine. On the basis of this finding, the present inventors have conducted further studies, and have found that when a drug preparation is formed from a composition which contains pitavastatin, a salt thereof, or an ester thereof and which initiates release thereof at least in the stomach and, in combination, an enteric composition which contains pitavastatin, a salt thereof, or an ester thereof, the drug preparation can attain an appropriate effective blood level thereof immediately after administration and can maintain such an appropriate blood level for a long period of time. The present invention has been accomplished on the basis of these findings.

Summary of Invention Paragraph:

[0008] Accordingly, the present invention provides a pharmaceutical composition characterized by comprising a composition (A) which contains pitavastatin, a salt thereof, or an ester thereof and which initiates release thereof at least in the stomach, and an enteric composition (B) which contains pitavastatin, a salt thereof, or an ester thereof.

Brief Description of Drawings Paragraph:

[0010] FIG. 2 shows the change in blood plasma level of pitavastatin in the case where pitavastatin was perorally administered in dogs, in the form of the compositions of Example 5 and Comparative Examples 1 and 2.

Detail Description Paragraph:

[0011] The characteristic feature of the pharmaceutical composition of the present invention is that the composition comprises a composition (A) which contains pitavastatin, a salt thereof, or an ester thereof (hereinafter these compounds are collectively referred to as pitavastatin) and which initiates release of pitavastatin at least in the stomach, and an enteric composition (B) which contains pitavastatin. Herein, the composition (A) initiates release of pitavastatin at least in the stomach. The composition (A) may release, in the stomach, the practical portion of pitavastatin or a portion of pitavastatin. However, a composition (A) which releases at least 30 mass % pitavastatin in the stomach is preferred, from the viewpoint of reliable attainment of an effective blood level of pitavastatin immediately after administration.

Detail Description Paragraph:

[0012] Preferably, the enteric composition of (B) is a composition prepared by coating pitavastatin or a pitavastatin-containing composition with a component which dissolves at a pH of 3.0 or higher. Alternatively, a composition prepared by mixing pitavastatin with a component which dissolves at a pH of 3.0 or higher is preferred for the enteric composition of (B). Accordingly, the portion of

Detail Description Paragraph:

Detail Description Paragraph:

Detail Description Paragraph:

Detail Description Paragraph:

http://jupiter2:9000/bin/cgi-bin/accum_query.pl?MODE=%20%20%20%20Display%20%... 4/18/2007

Detail Description Paragraph:

Detail Description Paragraph:

Detail Description Paragraph:

Detail Description Paragraph:

Detail Description Paragraph:

Detail Description Paragraph:

http://jupiter2:9000/bin/cgi-bin/accum_query.pl?MODE=%20%20%20%20Display%20%... 4/18/2007

http://jupiter2:9000/bin/cgi-bin/accum query.pl?MODE=%20%20%20%20Display%20%... 4/18/2007

the preferable ratio is within a range of 15:1 to 1:30, more preferably 10:1 to 1:20, particularly preferably 5:1 to 1:15. When the percentage of release is at least 30 mass % and less than 60 mass %, the preferable ratio is within a range of 30:1 to 1:20, more preferably 20:1 to 1:15, particularly preferably 15:1 to 1:10.

Detail Description Paragraph:

[0051] No particular limitation is imposed on the amount of pitavastatin incorporated into the controlled release pharmaceutical composition of the present invention, and the preferable amount is within a range of 0.01 to 60 mass %.

Detail Description Paragraph:

[0052] The amounts of ordinary bases, sustained release components, and enteric components incorporated into the controlled release pharmaceutical composition of the present invention vary depending on the types of these additives, design of release control of drugs, etc. However, the amount of each additive preferably is within a range of 0.01 to 80 mass %, more preferably 0.1 to 50 mass %, particularly preferably 1.0 to 30 mass %.

Detail Description Paragraph:

[0054] Examples of preferred drug forms of the controlled release pharmaceutical composition of the present invention for attaining a satisfactory blood level of pitavastatin include the following forms:

Detail Description Paragraph:

[0065] The daily dose of pitavastatin contained in the thus-obtained controlled release pharmaceutical composition of the present invention is adjusted to 0.5 to 64 mg, preferably 1 to 32 mg, particularly preferably 4 to 16 mg.

Detail Description Paragraph:

[0068] To purified water (3000.0 g), pitavastatin calcium salt (240.0 g), sucrose (615.0 g), hydroxypropylmethylcellulose (registered trade mark: TC-5R) (360.0 g), magnesium alumino metasilicate (48.0 g), talc (12.0 g) and triethyl citrate (45.0 g) were added, and these components were dissolved/dispersed by use of a disperser, to thereby prepare a drug-containing coating solution.

Detail Description Paragraph:

[0072] To purified water (1000.0 g), pitavastatin calcium salt (80.0 g), sucrose (205.0 g), hydroxypropylmethylcellulose (registered trade mark: TC-5R) (120.0 g), magnesium alumino metasilicate (16.0 g), talc (4.0 g) and triethyl citrate (15.0 g) were added, and these components were dissolved/dispersed by use of a disperser, to thereby prepare a drug-containing coating solution.

Detail Description Paragraph:

[0078] To purified water (500.0 g), pitavastatin calcium salt (40.0 g), sucrose (102.5 g), hydroxypropylmethylcellulose (registered trade mark: TC-5R) (60.0 g), magnesium alumino metasilicate (8.0 g), talc (2.0 g) and triethyl citrate (7.5 g) were added, and these components were dissolved/dispersed by use of a disperser, to thereby prepare a drug-containing coating solution.

Detail Description Paragraph:

[0079] To ethanol (1312.5 g), hydroxypropylmethylcellulose phthalate (161.7 g), triethyl citrate (16.1 g) and talc (32.2 g) were added, and these components were dissolved/dispersed by use of a disperser, to thereby prepare an enteric coating solution.

Detail Description Paragraph:

[0081] To purified water (500.0 g), pitavastatin calcium salt (40.0 g), sucrose (102.5 g), hydroxypropylmethylcellulose (registered trade mark: TC-5R) (60.0 g), magnesium alumino metasilicate (8.0 g), talc (2.0 g) and triethyl citrate (7.5 g) were added, and these components were dissolved/dispersed by use of a disperser, to

thereby prepare a drug-containing coating solution.

Detail Description Paragraph:

[0085] Hydroxypropylmethylcellulose (registered trade mark: TC-5R) (325.0 g) was dissolved in purified water (1500 g) by use of a disperser. Ethylcellulose (100.0 g), triethyl citrate (32.5 g) and talc (82.5 g) were dissolved/dispersed in ethanol (4000.0 g) by use of a disperser. The two solutions were mixed, to thereby prepare a coating solution.

Detail Description Paragraph:

[0092] A pitavastatin calcium salt (60.0 g), ethyl acrylate.methyl methacrylate (chlorotrimethylammonio)ethyl methacrylate copolymer (registered trade mark: Eudragit RS) (982.8 g), lactose (655.2 g), low-substituted hydroxypropylcellulose (90.0 g) and magnesium alumino metasilicate (12.0 g) were mixed, to thereby prepare a uniform powder mixture. A suitable amount of ethanol was added thereto, and the resultant mixture was granulated by a high-shear granulator, to thereby produce a granular product (1800.0 g).

Detail Description Paragraph:

[0093] To ethanol (5225.5 g), hydroxypropylmethylcellulose acetate succinate (311.85 g), triethyl citrate (31.05 g) and talc (62.1 g) were added, and these components were dissolved/dispersed by use of a disperser, to thereby prepare an enteric coating solution.

Detail Description Paragraph:

[0104] The granular product A (900.0 g), the enteric granular product (1170.0 g), crystalline cellulose (2146.2 g), hydroxypropylmethylcellulose (registered trade mark: TC-5R) (147.0 g), low-substituted hydroxypropylcellulose (588.0 g) and magnesium stearate (58.8 g) were mixed and compressed, to thereby produce 10,000 tablets, each tablet having a weight of 501.0 mg.

Detail Description Paragraph:

[0107] Pitavastatin calcium salt (40.0 g), lactose (3450.0 g), low-substituted hydroxypropylcellulose (770.0 g), hydroxypropylmethylcellulose (registered trade mark: TC-5R) (88.0 g) and magnesium alumino metasilicate (8.0 g) were mixed, to thereby prepare a uniform powder mixture. A suitable amount of purified water was added thereto, and the resultant mixture was granulated by a high-shear granulator. The thus-obtained granular product was mixed with magnesium stearate (44.0 g), to thereby produce a mixed granular product (4400.0 g).

Detail Description Paragraph:

[0112] A mixture containing the granular product A (600.0 g), crystalline cellulose (438.0 g), hydroxypropylmethylcellulose (registered trade mark: TC-5R) (30.0 g), low-substituted hydroxypropylcellulose (120.0 g) and magnesium stearate (12.0 g), and another mixture containing the enteric granular product A (780.0 g), crystalline cellulose (569.4 g), hydroxypropylmethylcellulose (registered trade mark: TC-5R) (39.0 g), low-substituted hydroxypropylcellulose (156.0 g) and magnesium stearate (15.6 g) were compressed, to thereby produce 10,000 two-layer tablets, each tablet having a weight of 276.0 mg (A layer: 120.0 mg, B layer: 156.0 mg).

Detail Description Paragraph:

[0115] Pitavastatin calcium salt (80.0 g), carboxymethylethylcellulose (1010.4 g), lactose (673.6 g), low-substituted hydroxypropylcellulose (200.0 g) and magnesium alumino metasilicate (16.0 g) were mixed, to thereby prepare a uniform powder mixture. A suitable amount of ethanol was added thereto, and the resultant mixture was granulated by a high-shear granulator. The thus-obtained granular product was mixed with magnesium stearate (20.0 g), to thereby produce an enteric mixed granular product (2000.0 g).

Detail Description Paragraph:

[0118] Pitavastatin calcium salt (80.0 g), ethyl acrylate.methyl methacrylate. (chlorotrimethylammonio)ethyl methacrylate copolymer (registered trade mark: Eudragit RS) (568.8 g), lactose (1603.2 g), low-substituted hydroxypropylcellulose (120.0 g) and magnesium alumino metasilicate (16.0 g) were mixed, to thereby prepare a uniform powder mixture. A suitable amount of ethanol was added thereto, and the resultant mixture was granulated by a high-shear granulator. The thus-obtained granular product was mixed with magnesium stearate (12.0 g), to thereby produce a mixed granular product (2400.0 g).

Detail Description Paragraph:

[0119] Pitavastatin calcium salt (40.0 g), carboxymethylethylcellulose (505.2 g), lactose (336.8 g), low-substituted hydroxypropylcellulose (100.0 g) and magnesium alumino metasilicate (8.0 g) were mixed, to thereby prepare a uniform powder mixture. A suitable amount of ethanol was added thereto, and the resultant mixture was granulated by a high-shear granulator. The thus-obtained granular product was mixed with magnesium stearate (10.0 g), to thereby produce an enteric mixed granular product (1000.0 g).

Detail Description Paragraph:

[0122] Pitavastatin calcium salt (60.0 g), lactose (324.0 g), low-substituted hydroxypropylcellulose (45.0 g), hydroxypropylmethylcellulose (registered trade mark: TC-5R) (7.5 g) and magnesium alumino metasilicate (9.0 g) were mixed, to thereby prepare a uniform powder mixture. A suitable amount of purified water was added thereto, and the resultant mixture was granulated by a high-shear granulator. The thus-obtained granular product and magnesium stearate (4.5 g) were mixed and compressed, to thereby produce 10,000 uncoated tablets (total 450.0 g), each tablet having a weight of 45.0 mg.

Detail Description Paragraph:

[0129] Pitavastatin calcium salt (90.0 g), methacrylate.methyl methacrylate copolymer (registered trade mark: Eudragit S) (380.7 g), lactose (253.8 g), low-substituted hydroxypropylcellulose (48.5 g) and magnesium alumino metasilicate (18.0 g) were mixed, to thereby prepare a uniform powder mixture. A suitable amount of ethanol was added thereto, and the resultant mixture was granulated by a high-shear granulator. The thus-obtained granular product and magnesium stearate (9.0 g) were mixed and compressed, to thereby produce 10,000 enteric tablets (800.0 g), each tablet having a weight of 80.0 mg.

Detail Description Paragraph:

[0132] To purified water (4000.0 g), pitavastatin calcium salt (120.0 g), sucrose (143.5 g), hydroxypropylmethylcellulose (registered trade mark: TC-5R) (84.0 g), magnesium alumino metasilicate (24.0 g), talc (6.0 g) and triethyl citrate (22.5 g) were added, and these components were dissolved/dispersed by use of a disperser, to thereby prepare a drug-containing coating solution.

Detail Description Paragraph:

[0136] The enteric granular product (1040.0 g), crystalline cellulose (759.2 g), hydroxypropylmethylcellulose (registered trade mark: TC-5R) (52.0 g), low-substituted hydroxypropylcellulose (208.0 g), and magnesium stearate (20.8 g) were mixed and compressed, to thereby produce 10,000 core tablets (2080.0 g), each tablet having a weight of 208.0 mg.

Detail Description Paragraph:

[0139] Pitavastatin-controlled-release tablets can be produced in accordance with the formulations shown in Table 15. Specifically, after preparation of core tablets, a drug-controlled-release coating layer is formed on the core tablets. Methacrylate.methyl methacrylate copolymer (trade name: Eudragit L, available from: Higuchi Inc.) is an enteric polymer which dissolves at a pH of 6 to 14, and ethylcellulose (trade name: Ethocel, product of Dow Chemical) is a sustained-

release polymer which is insoluble regardless of pH. The tablets obtained in Production Examples are pitavastatin-controlled-release tablets.

Detail Description Paragraph:

[0140] The tablets obtained in Production Example 1 are further coated with a pitavastatin-containing composition of Table 16. The tablet product of Example 14, which is formed of an outermost layer (immediately release portion) and an inner layer (controlled release portion) in combination, maintains reliable release of pitavastatin immediately after administration over a long period of time.

Detail Description Paragraph:

[0141] A pitavastatin-controlled-release granular product can be produced in accordance with the formulation shown in Table 17. Specifically, after preparation of core granules, a drug-controlled-release coating layer is formed on the core granules. The core granules and the drug-controlled-release-layer-coated core granules are mixed, to thereby yield a controlled release granular product.

Detail Description Paragraph:

[0143] A pitavastatin-controlled-release capsule product can be obtained by mixing core granules of Table 17; drug-controlled-release layer-coated granules of Table 17; and drug-controlled-release layer-coated granules of Table 18 and by charging the mixture in capsules.

Detail Description Paragraph:

[0144] A pitavastatin-controlled-release tablet product can be obtained by mixing core granules of Table 17; drug-controlled-release layer-coated granules of Table 17; and drug-controlled-release layer-coated granules of Table 18 and by compressed the mixture.

Detail Description Paragraph:

[0145] A pitavastatin-controlled-release multiple-layer tablet product can be obtained by individually compressing a layer of core granules of Table 17 and a layer of drug-controlled-release layer-coated granules of Table 17.

Detail Description Paragraph:

[0146] A pitavastatin-controlled-release tablet product can be produced in accordance with the formulation shown in Table 19. Specifically, after preparation of core tablets of Table 19, an outer layer component shown in Table 19 is formed on the core tablets by compressing, to thereby yield core-shell tablets.

Detail Description Paragraph:

[0149] Pitavastatin calcium salt (400.0 g), lactose (2160.0 g), low-substituted hydroxypropylcellulose (300.0 g), hydroxypropylmethylcellulose (registered trade mark: TC-5R) (50.0 g) and magnesium alumino metasilicate (60.0 g) were mixed, to thereby prepare a uniform powder mixture. Purified water (594.0 g) was added thereto, and the mixture was granulated by a high-shear granulator. The thus-obtained granular product and magnesium stearate (30.0 g) were mixed and compressed, to thereby produce 25,000 tablets, each tablet having a weight of 120.0 mg.

Detail Description Paragraph:

[0151] Pitavastatin calcium salt (320.0 g), lactose (960.0 g), hydroxypropylmethylcellulose (registered trade mark: Methocel K100LV) (1600.0 g), magnesium alumino metasilicate (128.0 g) and low-substituted hydroxypropylcellulose (160.0 g) were mixed, to thereby prepare a uniform powder mixture. Purified water (1108.8 g) was added thereto, and the resultant mixture was granulated by a high-shear granulator. The thus-obtained granular product and magnesium stearate (32.0 g) were mixed and compressed, to thereby produce 20,000 tablets, each tablet having a weight of 160.0 mg.

Detail Description Paragraph:

[0153] The dissolution property of pitavastatin from drug preparations; i.e., a drug preparation of Example 5 (composition (A) and enteric composition (B)); that of Comparative Example 1 (ordinary preparation); and that of Comparative Example 2 (sustained release preparation) was investigated in accordance with the following method.

Detail Description Paragraph:

[0157] Each of drug preparations of Example 5 and Comparative Examples 1 and 2 was added to the artificial gastric juice medium (900 mL) (pH: 1.2), and the medium was stirred at 37. \pm .1.degree. C. with a paddle rotation of 100 rpm for three hours after starting of the test. Immediately thereafter, the dissolution test medium was changed to the artificial intestinal juice medium (900 mL) (pH: 6.8), and the dissolution test was continued for 21 hours at 37. \pm .1.degree. C. with a paddle rotation of 100 rpm. The amount of dissolved drug was determined. A sample solution sampled at each timing was filtered by means of a membrane filter made of cellulose acetate (pore size: 0.45 μ m, DISMIC-25cs, product of Toyo Roshi), and the percent of dissolution of pitavastatin was determined with high performance liquid chromatography by use of a reverse-phase column (Develosil ODS-HG-5, product of Nomura Chemical). The results are shown in FIG. 1.

Detail Description Paragraph:

[0159] Each of the drug preparations prepared in Example 5, Comparative Example 1, and Comparative Example 2 was orally administered to an HRA beagle dog (body weight: about 10 kg) under fasting. After administration, blood was collected at different points in time for 24 hours, and the collected blood was subjected to centrifugation, whereby blood pitavastatin level was determined with high performance liquid chromatography. The results are shown in Table 22 and FIG. 2. The pharmacokinetic values are shown in Table 23. The results show that the controlled release drug preparation of the present invention reduces C.sub.max by 50% as compared with the ordinary drug preparation, while providing equivalent AUC comparable to that of the ordinary drug preparation. By contrast, in the case of the sustained release drug with zero-order release, C.sub.max and AUC are both significantly reduced as compared with the ordinary drug preparation. Therefore, the controlled release drug preparation of the present invention was found to be safe and highly effective.

Detail Description Paragraph:

[0161] By use of the controlled release pharmaceutical composition of the present invention, the blood level of pitavastatin can be maintained at an appropriate level immediately after administration over a long period of time. Thus, highly safety and effective treatment of hypercholesterolemia can be performed.

Detail Description Table CWU:

1TABLE 1 Absorption sites of pitavastatin in rat (in situ loop test) Absorption percentage (%) Sites 0.5 h 1 h Stomach 8.6 \pm 1.1 14.4 \pm 1.6 Duodenum 65.9 \pm 2.1 60.3 \pm 10.2 Small 23.1 \pm 3.8 21.3 \pm 3.0 intestine Large 34.4 \pm 3.7 51.6 \pm 6.7 intestine

Detail Description Table CWU:

2TABLE 2 Formulation Components (mg) Enteric Granular Purified sucrose spheres 114.00 granular product A Pitavastatin calcium salt 12.00 product A Sucrose 30.75 Hydroxypropylmethylcellulose 18.00 (Registered trade mark: TC-5R) Magnesium alumino metasilicate 2.40 Talc 0.60 Triethyl citrate 2.25 Subtotal 180.00 Methacrylate .multidot. methyl methacrylate 41.58 copolymer (Registered trade mark: Eudragit L) Talc 8.28 Triethyl citrate 4.14 Subtotal 234.00 Pitavastatin calcium salt 4.00 Sucrose 10.25 Hydroxypropylmethylcellulos- e (Registered trade 6.00 mark: TC-5R) Magnesium alumino metasilicate 0.80 Talc 0.20 Triethyl citrate 0.75 Total 256.00

Detail Description Table CWU:

3TABLE 3 Formulation Components (mg) Enteric Granular Granular product A 30.00 granular product Ethyl acrylate .multidot. methyl 12.40 product methacrylate .multidot. (chloro- trimethylammonio) ethyl methacrylate copolymer (Registered trade mark: Eudragit RS) Hydroxypropylcellulose 3.10 Talc 0.95 Triethyl citrate 1.55 Subtotal 48.00 Pitavastatin calcium salt 4.00 Sucrose 10.25 Hydroxypropylmethylcellulose (Registered 6.00 trade mark: TC-5R) Magnesium alumino metasilicate 0.80 Talc 0.20 Triethyl citrate 0.75 Hydroxypropylmethylcellulose phthalate 16.17 Talc 3.22 Triethyl citrate 1.61 Subtotal 91.00 Pitavastatin calcium salt 4.00 Sucrose 10.25 Hydroxypropylmethylcellulose (Registered trade 6.00 mark: TC-5R) Magnesium alumino metasilicate 0.80 Talc 0.20 Triethyl citrate 0.75 Total 113.00

Detail Description Table CWU:

4 TABLE 4 Formulation Components (mg) Granular Granular product A 180.00 product Ethylcellulose 10.00 Hydroxypropylmethylcellulose 32.50 (Registered trade mark: TC-5R) Talc 8.25 Triethyl citrate 3.25 Subtotal 234.00 Enteric Granular product A 60.00 granular Methacrylate .multidot. methyl methacrylate 13.86 product copolymer (Registered trade mark: Eudragit S) Talc 2.76 Triethyl citrate 1.38 Subtotal 78.00 Total 312.00

Detail Description Table CWU:

5TABLE 5 Formulation Components (mg) Granular product A 60.00 Granular Pitavastatin calcium salt 4.00 product Ethyl acrylate .multidot. methyl 65.52 methacrylate. (chlorotrimethylammonio)ethyl methacrylate copolymer (Registered trade mark: Eudragit RS) Lactose 43.68 Low-substituted hydroxypropylcellulose 6.00 Magnesium alumino metasilicate 0.80 Subtotal 120.00 Enteric Granular product A 90.00 granular Hydroxypropylmethylcellulose acetate 20.79 product succinate Talc 4.14 Triethyl citrate 2.07 Subtotal 117.00 Total 297.00

Detail Description Table CWU:

7TABLE 7 Formulation Components (mg) Granular product A 90.00 Enteric Granular product A 90.00 granular Methacrylate .multidot. ethyl acrylate copolymer 20.79 product (Registered trade mark: Eudragit L30D-55) Talc 4.14 Triethyl citrate 2.07 Subtotal 117.00 Crystalline cellulose 214.62 Hydroxypropylmethylcellulose 14.70 (Registered trade mark: TC-5R) Low-substituted hydroxypropylcellulose 58.80 Magnesium stearate 5.88 Total 501.00

Detail Description Table CWU:

8TABLE 8 Formulation Components (mg) Enteric granular product A 195.00 Mixed Pitavastatin calcium salt 4.00 granular Lactose 345.00 product Low-substituted hydroxypropylcellulose 77.00 Hydroxypropylmethylcellulose 8.80 (Registered trade mark: TC-5R) Magnesium alumino metasilicate 0.80 Magnesium stearate 4.40 Subtotal 440.00 Total 635.00

Detail Description Table CWU:

9TABLE 9 Formulation Components (mg) A layer Granular product A 60.00 Crystalline cellulose 43.80 Hydroxypropylmethyl cellulose 3.00 (Registered trade mark: TC-5R) Low-substituted hydroxypropylcellulose 12.00 Magnesium stearate 1.20 Subtotal 120.00 B layer Enteric granular product A 78.00 Crystalline cellulose 56.94 Hydroxypropylmethylcellulose 3.90 (Registered trade mark: TC-5R) Low-substituted hydroxypropylcellulose 15.60 Magnesium stearate 1.56 Subtotal 156.00 Total 276.00

Detail Description Table CWU:

10TABLE 10 Formulation Components (mg) A layer Mixed granular product 440.00 B layer Pitavastatin calcium salt 8.00 Carboxymethylethylcellulose 101.04 Lactose 67.36 Low-substituted hydroxypropylcellulose 20.00 Magnesium alumino metasilicate 1.60 Magnesium stearate 2.00 Subtotal 200.00 Total 640.00

Detail Description Table CWU:

11TABLE 11 Formulation Components (mg) A layer Pitavastatin calcium salt 8.00 Ethyl acrylate .multidot. methyl methac- 56.88 rylate .multidot. (chlorotrimethylammonio) ethyl methacrylate copolymer (Registered trade mark: Eudragit RS) Lactose 160.32 Low-substituted hydroxypropylcellulose 12.00 Magnesium alumino metasilicate 1.60 Magnesium stearate 1.20 Subtotal 240.00 B layer Pitavastatin calcium salt 4.00 Carboxymethylethylcellulose 50.52 Lactose 33.68 Low-substituted hydroxypropylcellulose 10.00 Magnesium alumino metasilicate 0.80 Magnesium stearate 1.00 Subtotal 100.00 Total 340.00

Detail Description Table CWU:

12TABLE 12 Formulation Components (mg) Core Pitavastatin calcium salt 6.00 tablet Lactose 32.40 Low-substituted hydroxypropylcellulose 4.50 Hydroxypropylmethylcellulose 0.75 (Registered trade mark: TC-5R) Magnesium alumino metasilicate 0.90 Magnesium stearate 0.45 Methacrylate .multidot. methyl methacrylate copolymer 15.40 (Registered trade mark: Eudragit S) Talc 1.54 Triethyl citrate 3.06 Subtotal 65.00 Outer Mixed granular product 220.00 layer Total 285.00

Detail Description Table CWU:

13TABLE 13 Formulation Components (mg) Core Pitavastatin calcium salt 9.00 tablet Methacrylate .multidot. methyl methacrylate copolymer 38.07 (Registered trade mark: Eudragit S) Lactose 25.38 Low-substituted hydroxypropylcellulose 4.85 Magnesium alumino metasilicate 1.80 Magnesium stearate 0.90 Subtotal 80.00 Outer Mixed granular product 330.00 layer Total 410.00

Detail Description Table CWU:

14TABLE 14 Formulation Components (mg) Core Enteric Purified sucrose spheres 40.00 tablet granular Pitavastatin calcium salt 12.00 product Sucrose 14.35 Hydroxypropylmethylcellulose 8.40 (Registered trade mark: TC-5R) Magnesium alumino metasilicate 2.40 Talc 0.60 Triethyl citrate 2.25 Cellulose acetate phthalate 18.48 Talc 3.68 Triethyl citrate 1.84 Subtotal 104.00 Crystalline cellulose 75.92 Hydroxypropylmethylcellulose (Registered 5.20 trade mark: TC-5R) Low-substituted hydroxypropylcellulose 20.80 Magnesium stearate 2.08 Subtotal 208.00 Outer Mixed granular product 440.00 layer Total 648.00

Detail Description Table CWU:

15 TABLE 15 Production Examples (mg/T) Components 1 2 Core tablet Pitavastatin calcium salt 8.0 8.0 Lactose 94.4 94.4 Low-substituted hydroxypropylcel- 12.0 12.0 lulose Hydroxypropylmethylcellulose 2.0 2.0 Magnesium alumino metasilicate 2.4 2.4 Magnesium stearate 1.2 1.2 Subtotal 120.0 120.0 Drug-controlled- Methacrylate .multidot. methyl methacrylate 8.0 -- release coating copolymer layer Ethylcellulose -- 8.0 Triethyl citrate 1.2 1.2 Talc 2.5 2.5 Titanium oxide 1.3 1.3 Subtotal 13.0 13.0 Total 133.0 133.0

Detail Description Table CWU:

16 TABLE 16 Example 14 Components (mg/T) Inner Tablets (Production Example 1) 133.0 layer Outermost Pitavastatin calcium salt 8.0 layer Lactose 304.0 Low-substituted 36.0 hydroxypropylcellulose Hydroxypropylmethylcellulose 6.0 Magnesium alumino metasilicate 2.4 Magnesium stearate 3.6 Subtotal 360.0 Total 493.0

Detail Description Table CWU:

17 TABLE 17 Example 15 Components (mg/day) Core Pitavastatin calcium salt 8.0 granular D-mannitol 334.0 product Calcium Carboxymethylcellulose 21.0 Hydroxypropylcellulose 21.0 Magnesium alumino metasilicate 16.0 Subtotal 400.0 Drug- Methacrylate.methyl methacrylate 54.0 controlled- copolymer release Triethyl citrate 12.0 coating Talc 50.0 layer Titanium oxide 4.0 Subtotal 120.0

Detail Description Table CWU:

18 TABLE 18 Example 16 Components (mg/day) Core Pitavastatin calcium salt 8.0 granular D-mannitol 334.0 product Calcium Carboxymethylcellulose 21.0 Hydroxypropylcellulose 21.0 Magnesium alumino metasilicate 16.0 Subtotal 400.0

Drug- Methacrylate.methyl methacrylate 54.0 controlled- copolymer release
 Ethylcellulose 27.0 coating Triethyl citrate 18.0 layer Talc 75.0 Titanium oxide
 6.0 Subtotal 180.0

Detail Description Table CWU:

19 TABLE 19 Production Example 3 Components (mg/T) Core Pitavastatin calcium salt
 8.0 tablet Lactose 46.4 Crospovidone 20.0 Hydroxypropylmethylcellulose 2.0
 Magnesium alumino metasilicate 2.4 Ethyl acrylate.methyl methacrylate. 20.0
 (chlorotrimethylammonio)ethyl methacrylate copolymer*.sup.1 Monoglyceride 20.0
 Magnesium stearate 1.2 Total 120.0 Outer Pitavastatin calcium salt 8.0 layer
 Lactose 304.0 Low-substituted 36.0 hydroxypropylcellulose
Hydroxypropylmethylcellulose 6.0 Magnesium alumino metasilicate 2.4 Magnesium
 stearate 3.6 Subtotal 360.0 *.sup.1Eudragit RS sold by Higuchi Inc.

Detail Description Table CWU:

20 TABLE 20 Components Formulation (mg) Pitavastatin calcium salt 16.00 Lactose
 86.40 Low-substituted hydroxypropylcellulose 12.00 Hydroxypropylmethylcellulos 2.00
 (Registered trade mark: TC-5R) Magnesium alumino metasilicate 2.40 Magnesium
 stearate 1.20 Total 120.00

Detail Description Table CWU:

21 TABLE 21 Components Formulation (mg) Pitavastatin calcium salt 16.00 Lactose
 48.00 Hydroxypropylmethylcellulose 80.00 (Registered trade mark: Methocel K100LV)
 Magnesium alumino metasilicate 6.40 Low-substituted hydroxypropylcellulose 8.00
 Magnesium stearate 1.60 Total 160.00

Detail Description Table CWU:

22TABLE 22 Pitavastatin blood plasma level at each sampling time of the composition
 prepared in Example 5 and Comparative Examples 1 and 2 in beagle dog oral
 administration Time (h) 0 0.5 1 1.5 2 3 4 6 8 10 24 Ex. 5 0 229.7 505.7 533.3 536.3
 386.0 373.0 229.3 112.5 59.8 8.2 (ng/mL) Comp. Ex. 1 0 1246.3 1179.7 690.8 475.2
 323.7 226.5 155.8 49.3 29.0 8.3 (ng/mL) Comp. Ex. 2 0 10.2 52.2 73.7 76.5 131.3
 161.8 171.7 49.0 25.3 3.2 (ng/mL)

CLAIMS:

1. A controlled release pharmaceutical composition characterized by comprising a composition (A) which contains pitavastatin, a salt thereof, or an ester thereof and which initiates release thereof at least in the stomach, and an enteric composition (B) which contains pitavastatin, a salt thereof, or an ester thereof.
2. A controlled release pharmaceutical composition as described in claim 1, wherein the composition (A) releases in the stomach at least 30 mass % pitavastatin, a salt thereof, or an ester thereof contained in the composition (A).
3. A controlled release pharmaceutical composition as described in claim 1, which has a ratio by mass of pitavastatin, a salt thereof, or an ester thereof contained in the composition (A) to pitavastatin, a salt thereof, or an ester thereof contained in the enteric composition (B) falling within a range of 30:1 to 1:40.
4. A controlled release pharmaceutical composition as described in claim 1, wherein the composition (A) releases in the stomach at least 85 mass % pitavastatin, a salt thereof, or an ester thereof contained in the composition (A), and the controlled release pharmaceutical composition has a ratio by mass of pitavastatin, a salt thereof, or an ester thereof contained in the composition (A) to pitavastatin, a salt thereof, or an ester thereof contained in the enteric composition (B) falling within a range of 1:1 to 1:40.
5. A controlled release pharmaceutical composition as described in claim 1, wherein the composition (A) releases in the stomach at least 60 mass % and less than 85

mass % pitavastatin, a salt thereof, or an ester thereof contained in the composition (A), and the controlled release pharmaceutical composition has a ratio by mass of pitavastatin, a salt thereof, or an ester thereof contained in the composition (A) to pitavastatin, a salt thereof, or an ester thereof contained in the enteric composition (B) falling within a range of 15:1 to 1:30.

6. A controlled release pharmaceutical composition as described in claim 1, wherein the composition (A) releases in the stomach at least 30 mass % and less than 60 mass % pitavastatin, a salt thereof, or an ester thereof contained in the composition (B), and the controlled release pharmaceutical composition has a ratio by mass of pitavastatin, a salt thereof, or an ester thereof contained in the composition (A) to pitavastatin, a salt thereof, or an ester thereof contained in the enteric composition (B) falling within a range of 30:1 to 1:20.

7. A controlled release pharmaceutical composition as described in any one of claims 1 to 6, wherein the enteric composition (B) is obtained by coating pitavastatin, a salt thereof, or an ester thereof or a composition containing the same with a component which dissolves at a pH of 3 or higher or by mixing pitavastatin, a salt thereof, or an ester thereof with a component which dissolves at a pH of 3 or higher.

[Previous Doc](#)

[Next Doc](#)

[Go to Doc#](#)